

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



King Saud University
Saudi Journal of Biological Sciences

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Optimization of multiple shoot induction and plant regeneration in Indian barley (*Hordeum vulgare*) cultivars using mature embryos



Hassan Rostami ^a, Archana Giri ^a, Amir Sasan Mozaffari Nejad ^{b,*}, Amir Moslem ^c

^a Center for Biotechnology, Institute of Science and Technology, Jawaharlal Nehru Technological University, Hyderabad, Andhra Pradesh, India

^b Member of Young Researchers and Elite Club, Garmsar Branch, Islamic Azad University, Garmsar, Iran

^c Department of Agronomy and Plant Breeding, University of Tehran, College of Agriculture and Natural Resources, Karaj 4111, Iran

Received 6 January 2013; revised 14 February 2013; accepted 15 February 2013

Available online 14 March 2013

KEYWORDS

Hordeum vulgare;
Direct organogenesis;
Plant regeneration;
Mature embryos

Abstract Barley is the fourth most important crop in the world. Development of a regeneration system using immature embryos is both time consuming and laborious. The present study was initiated with a view to develop a regeneration system in six genotypes of Indian barley (*Hordeum vulgare*) cultivars as a prerequisite to transformation. The mature embryos were excised from seeds and cultured on MS medium supplemented with high and low concentrations of cytokinins and auxins respectively. The MS medium containing 3 mg/L N⁶-benzylaminopurine (BA) and 0.5 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D) was found to be the most effective for multiple shoot formation in HOR7231 cultivar that could produce 12 shoots per explant. The other cultivars HOR4409 and HOR3844 produced a minimum number of adventitious shoots (1.33 and 1.67 respectively) on MS medium supplemented with 1 mg/L BA and 0.3 mg/L 2,4-D. The elongated shoots were separated and successfully rooted on MS medium containing 1 mg/L indole-3-acetic acid (IAA). The response of different barley cultivars was found to be varying with respect to multiple shoot production. This is the first report of multiple shoot induction and plantlet regeneration in Indian cultivar of barley which would be useful for genetic transformation.

© 2013 King Saud University. Production and hosting by Elsevier B.V. All rights reserved.

* Corresponding author. Mobile: +91 888 545 9181.

E-mail address: asmozafarnejad@yahoo.in (A.S.M. Nejad).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

Barley is an annual cereal grain crop that comes from the family of *Poaceae* and is the fourth most important crop in the world after wheat, maize and rice (faostat.org). It is consumed as a major food and feed for animals as well as in the brewing industry (Ritala et al., 1993). Barley seeds are unique as they accumulate up to 15% protein by dry weight and be stored for a period as long as 10 years (Tanasienko et al.,

2011) and (Ritala et al., 2008). Considering its importance, there is a requirement for development of transgenic barley with new properties (Wan and Lemaux, 1994). Existence of high potent in vitro regeneration system is a prerequisite for efficient genetic transformation (Havrlentová et al., 2001). The regeneration system is supposed to be fast growing with stable plantlets suitable for hardening (Sood et al., 2011). Barley is one of the most difficult crops for in vitro tissue culture as it is a well known recalcitrant species like other monocots. Several explants of barley have been considered for their regeneration and transformation capacity and successful attempts have been reported including immature embryos (Wan and Lemaux, 1994), microspores (Jahne et al., 1994), protoplasts (Funatsuki et al., 1995), shoot meristematic cultures (Zhang et al., 1999), ovules (Holme et al., 2006). Other explants which were used for in vitro regeneration of cereal plants include: young leaves and leaf tissues (Saalbach and Koblitz, 1978) and (Lu et al., 1983), anthers (Haydu and Vasil, 1981), immature inflorescences (Havrlentová et al., 2001) and (Desai et al., 2004), nodal explants (Sharma et al., 2007), leaf base and apical meristems (Ganeshan et al., 2003) and (Ramesh et al., 2009).

Most regeneration protocols for barley are based on immature embryos in controlled conditions and the procedure for generation of donor plants is time consuming. Moreover, reproducible in vitro regeneration system even with immature embryos has mostly been restricted to a few cultivars (Li et al., 2009) like Golden Promise that is not commercially valuable. Nearly all the commercially important cultivars are poor in regeneration (Ganeshan et al., 2003) and have not been explored for in vitro studies (Chauhan and Kothari, 2004). On the other hand, explants like mature embryos and leaf base/apical meristems have several advantages over immature tissues as they are easier to handle, available throughout the year in a large quantity without any problems of seasonal effect on regeneration (Sharma et al., 2005a).

Direct regeneration of plants from explants is a faster and a time saving approach for obtaining whole plants without the callus interphase that can increase somaclonal variation (Zapata et al., 1999). Direct shoot induction and regeneration have been reported for some important cereals including oat (Zhang et al., 1996), maize (Zhang et al., 2002), sorghum (Baskaran and Jayabalan, 2005), barley (Sharma et al., 2004) and wheat (Sharma et al., 2005b). As the regeneration ability varies between genotypes, it is essential to screen different varieties for the response to in vitro culturing as it is a key step for genetic manipulation.

The objective of the present study was to investigate the regeneration capacity of barley cultivars originating from India using mature embryos as explant by direct shoot induction in order to use them for future molecular breeding programs.

2. Materials and methods

2.1. Preparation of explants

Barley seeds of six Indian varieties were provided by Gene Bank Department of Leibniz Institute of Plant Genetics and Cultivated Plant Research (IPK) – Germany. The accession numbers are shown in Table 1.

Seeds were dehusked and after washing with tap water surface sterilized by soaking in 70% ethanol for 1 min, rinsing with sterile distilled water, soaking in 0.1% (w/v) mercuric chloride solution for 7 min followed by rinsing with sterile distilled water (10 times). The seeds were kept in 250 ml conical flasks containing sterile water for 24 h at 4 °C. Mature embryos were excised from seeds by a scalpel and cultured on media.

2.2. Media composition, preparation and establishment of cultures

The seeds were sterilized after dehusking as described above and were incubated on direct shoot induction media (DSIM) for induction of multiple shoots. DSIM was MS basal media by Murashige and Skoog (1962) supplemented with high level of BA and low concentration of 2,4-D as shown in Table 1.

The cultures were kept in light at 22 ± 1 °C in 16/8 h of light/dark cycle regime with subculturing one in two weeks. Counting of multiple shoots was done for each explant after it was subcultured twice. The main shoots were removed during subculturing. Shoots which emerged from mature embryos directly without callus interphase were separated and transferred on rooting media. The regenerated plantlets were transferred on MS media containing 1 mg/L IAA for rooting.

2.3. Statistical analysis

In the direct shoot induction study the number of shoots per explants and efficiency of regeneration were recorded and ANOVA was conducted to investigate differences in above the parameters. All Statistical analysis was carried out by using DSAASTAT (available at: <http://www.unipg.it/~onofri/DAASTAT/DAASTAT.htm>).

3. Results and discussion

Mature embryos of six barley varieties were excised from 24 h imbibed surface sterilized seeds as initiation explants for direct embryogenesis on six different media containing different concentrations of BA and 2,4-D. Mature embryos started to germinate few days after culturing. The main shoots were cut to promote adventitious shoot initiation within the first week in order to retard the growth of primary shoots (Fig. 1A and

Table 1 Barley genotypes and combination of different media used in this study.

Cultivars used	HOR 3844, HOR 3272, HOR 4505, HOR 7231, HOR 4409, HOR 3838					
Levels of BAP and 24D in mg/L	1B + 0.3 D (1 mg/L BAP + 0.3 mg/L 2,4-D)	1B + 0.5D D (1 mg/L BAP + 0.5 mg/L 2,4-D)	2B + 0.3D D (2 mg/L BAP + 0.3 mg/L 2,4-D)	2B + 0.5D D (2 mg/L BAP + 0.5 mg/L 2,4-D)	3B + 0.3D D (3 mg/L BAPP + 0.3 mg/L 2,4-D)	3B + 0.5D D (3 mg/L BAP + 0.5 mg/L 2,4-D)




Figure 1 Different stages of induction direct multiple shoots from mature embryos of barley; A: Cutting the germinated embryos and initiation of shoot bud, B: Somatic embryo formation and development from shoot buds, C: Multiple shoots regeneration, D: Separating multiple shoots from each explants and E: Root induction of regenerated plants.

B). Subculturing the regenerated plantlets every two weeks led to multiplication and elongation of multiple shoots (Fig. 1C). Elongated shoots (2–3 cm length) from original explants were separated and subcultured on same media that could grow individually without albinism (Fig. 1D). All regenerated plantlets started rooting after two weeks from transferring them on MS media fortified with 1 mg/L IAA (Fig. 1E).

All varieties used in this study were able to induce adventitious shoots on six media. The range of multiple shoots varied from 1.33 for HOR4409 to 12 for HOR7231 (Fig. 2 and Table 2).

Best results were obtained with MS media containing 3 mg/L BA and 0.5 mg/L 2,4-D for all the varieties except of HOR3838. With HOR3838 the same result was obtained either on media containing 3 mg/L BA and 0.5 mg/L 2,4-D or 3 mg/L BA and 0.3 mg/L 2,4-D. The varieties HOR3844 and

HOR4409 had shown minimum response on media containing 1–2 mg/L BA and 0.3–0.5 mg/L 2,4-D. As it is shown in Table 2, all varieties exhibited the best response of induction and production of multiple adventitious shoots on MS media fortified with 3 mg/L BA and 0.5 mg/L 2,4-D. So it can be concluded that media enriched with high level of BA and low concentration of 2,4-D can be used to optimize media combination for regeneration and transformation of barley using mature embryo explants.

Thidiazuron (TDZ) was shown to have a positive influence on tissue culture of monocots (Schulze, 2007). TDZ was also used on cultivars HOR7231 and HOR3272 as these cultivars had shown good response with respect to induction of adventitious shoots. Substitution of BA with TDZ did not show any increase in the number of shoots. Similar results were reported by Ganeshan et al. (2006) with respect to barley. Ganeshan et al. (2003) reported a favorable response with leaf bases/apical meristems of barley genotypes to multiple shoot induction on two media containing TDZ and 2,4-D similar to those used by Gupta and Conger (1998) for switch grass. However the calli induced from leaf bases or scutella did not survive on TDZ containing media which is in contradiction to the report by Shan et al. (2000). The reasons for varying response could be the source of explants and the different requirement of phytohormones. Also the levels of TDZ used and the duration of incubation could induce different multiple shooting responses.

In the present study an efficient, fast and a reproducible in vitro regeneration system for Indian varieties of barley was developed using mature embryos. The results obtained (12 shoots per explants) in our experiment are comparable with those obtained for cultivar like Golden promise (9.5 shoots per explant) (Sharma et al., 2007).




Figure 2 Influence of different hormone combinations on direct shoot induction in barley genotypes.

Table 2 Minimum and maximum adventitious shoots induced with different hormone combination.

Varieties name	Minimum shoots induced	Media combination	Maximum shoot induced	Media combination
HOR 3844	1.67	1B + 0.3D	3.33	3B + 0.3D, 3B + 0.5D
HOR 3272	4.67	1B + 0.3D	10.00	3B + 0.5D
HOR 4505	3.33	1B + 0.3D	7.33	3B + 0.5D
HOR 7231	5.00	1B + 0.5D	12.00	3B + 0.5D
HOR 4409	1.33	1B + 0.3D	8.67	3B + 0.5D
HOR 3838	2.33	1B + 0.3D, 1B + 0.5D	4.67	3B + 0.5D

Most regeneration protocols in barley involve the use of immature embryos that can be obtained only after three months of cultivation of barley in green house followed by tissue culture and regeneration for two months. However, the protocol reported in the present paper involves regeneration of plantlets from mature embryos obtained from seeds within two months.

4. Conclusion

A protocol for tissue culture and regeneration of plantlets from mature embryos of Indian cultivars of barley has been developed. This protocol does not require green house as mature embryos obtained from seeds are used as explants. Significant differences have been observed between different cultivars and combinations of phytohormones. The results also indicate that although regeneration is possible with all cultivars, each cultivar requires a higher degree of standardization for improving the regeneration efficiency. Efficient regeneration protocol of this recalcitrant crop would aid in the development of protocols for genetic manipulation.

References

- Bhaskaran, P., Jayabalan, N., 2005. In vitro plant regeneration and mass propagation system for Sorghum bicolor-a valuable major cereal crop. *J. Agri. Technol.* 1, 345–363.
- Chauhan, M., Kothari, S.L., 2004. Optimization of nutrient levels in the medium increases the efficiency of callus induction and plant regeneration in recalcitrant Indian barley (*Hordeum vulgare* L.) in vitro. *In Vitro Cell. Dev.-Pl.* 40, 520–527.
- Desai, N.S., Suprasanna, P., Bapat, V.A., 2004. Simple and reproducible protocol for direct somatic embryogenesis from cultured immature inflorescence segments of sugarcane (*Saccharum* spp.). *Curr. Sci.* 87, 764–768.
- Funatsuki, H., Kuroda, H., Kihara, M., Lazzeri, P.A., Müller, E., Lörz, H., Kishinami, I., 1995. Fertile transgenic barley generated by direct DNA transfer to protoplasts. *Theor. Appl. Genet.* 91, 707–712.
- Ganeshan, S., Båga, M., Harvey, B.L., Rossnagel, B.G., Scoles, G.J., Chibbar, R.N., 2003. Production of multiple shoots from thidiazuron-treated mature embryos and leaf-base/apical meristems of barley (*Hordeum vulgare*). *Plant Cell Tiss. Org.* 73, 57–64.
- Ganeshan, S., Chodaparambil, S.V., Båga, M., Fowler, D.B., Hucl, P., Rossnagel, B.G., Chibbar, R.N., 2006. In vitro regeneration of cereals based on multiple shoot induction from mature embryos in response to thidiazuron. *Plant Cell Tiss. Org.* 85, 63–73.
- Gupta, S.D., Conger, B.V., 1998. In vitro differentiation of multiple shoot clumps from intact seedlings of switchgrass. *In Vitro Cell. Dev. Biol.-Plant.* 34, 196–202.
- Havrleňová, M., Faragó, J., Nešťáková, M., 2001. Regeneration of immature inflorescences of barley in vitro. *Biol. Plantarum* 44, 157–159.
- Haydu, Z., Vasil, I.K., 1981. Somatic embryogenesis and plant regeneration from leaf tissues and anthers of *Pennisetum purpureum* Schum. *Theor. Appl. Genet.* 59, 269–273.
- Holme, I.B., Brinch-Pedersen, H., Lange, M., Holm, P.B., 2006. Transformation of barley (*Hordeum vulgare* L.) by *Agrobacterium tumefaciens* infection of in vitro cultured ovules. *Plant Cell Rep.* 25, 1325–1335.
- Li, H.-P., Huang, T., Wang, C.-X., Liao, Y.-C., 2009. An efficient regeneration system of barley cultivars from leaf base segments. *Biol. Plantarum* 53, 733–736.
- Lu, C., Vasil, V., Vasil, I.K., 1983. Improved efficiency of somatic embryogenesis and plant regeneration in tissue cultures of maize (*Zea mays* L.). *Theor. Appl. Genet.* 66, 285–289.
- Murashige, T., Skoog, F., 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. plantarum* 15, 473–497.
- Ramesh, M., Murugiah, V., Gupta, A.K., 2009. Efficient in vitro plant regeneration via leaf base segments of indica rice (*Oryza sativa* L.). *Indian J. Exp. Biol.* 47, 68–74.
- Ritala, A., Mannonen, L., Aspegren, K., Salmenkallio-Marttila, M., Kurten, U., Hannus, R., Mendez Lozano, J., Teeri, T.H., Kauppinen, V., 1993. Stable transformation of barley tissue culture by particle bombardment. *Plant Cell Rep.* 12, 435–440.
- Ritala, A., Wahlström, E.H., Holkeri, H., Hafren, A., Mäkeläinen, K., Baez, J., Mäkinen, K., Nuutila, A.M., 2008. Production of a recombinant industrial protein using barley cell cultures. *Protein Express. Purif.* 59, 274–281.
- Saalbach, G., Koblit, H., 1978. Attempts to initiate callus formation from barley leaves. *Plant Sci. Lett.* 13, 165–169.
- Schulze, J., 2007. Improvements in cereal tissue culture by thidiazuron: a review. *Fruit, Veg. Cereal Sci. Biotech.* 1, 64–79.
- Shan, X., Li, D., Qu, R., 2000. Thidiazuron promotes in vitro regeneration of wheat and barley. *In Vitro Cell. Dev. Biol.-Plant.* 36, 207–210.
- Sharma, V.K., Hänsch, R., Mendel, R.R., Schulze, J., 2004. A highly efficient plant regeneration system through multiple shoot differentiation from commercial cultivars of barley (*Hordeum vulgare* L.) using meristematic shoot segments excised from germinated mature embryos. *Plant Cell Rep.* 23, 9–16.
- Sharma, V.K., Hänsch, R., Mendel, R.R., Schulze, J., 2005a. Influence of picloram and thidiazuron on high frequency plant regeneration in elite cultivars of wheat with long-term retention of morphogenicity using meristematic shoot segments. *Plant Breeding* 124, 242–246.
- Sharma, V.K., Hänsch, R., Mendel, R.R., Schulze, J., 2005b. Mature embryo axis-based high frequency somatic embryogenesis and plant regeneration from multiple cultivars of barley (*Hordeum vulgare* L.). *J. Exp. Bot.* 56, 1913–1922.
- Sharma, V.K., Hänsch, R., Mendel, R.R., Schulze, J., 2007. Node-derived cultures with high-morphogenic competence in barley and wheat. *Plant Cell Tiss. Org.* 88, 21–33.

- Sood, P., Bhattacharya, A., Sood, A., 2011. Problems and possibilities of monocot transformation. *Biol. Plantarum* 55, 1–15.
- Tanasienko, I.V., Yemets, A.I., Pirko, Y.V., Korhkovyy, V.I., Abumhadi, N., Blume, Ya.B., 2011. Generation of transgenic barley lines producing human lactoferrin using mutant alpha-tubulin gene as the selective marker. *Cytol. Genet.* 45, 1–6.
- Wan, Y., Lemaux, P.G., 1994. Generation of large numbers of independently transformed fertile barley plants. *Plant Physiol.* 104, 37–48.
- Zapata, C., Srivatanakul, M., Park, S.-H., Lee, B.-M., Salas, M.G., Smith, R.H., 1999. Improvements in shoot apex regeneration of two fiber crops: cotton and kenaf. *Plant Cell Tiss. Org.* 56, 185–191.
- Zhang, S., Zhang, H., Zhang, M.B., 1996. Production of Multiple Shoots from Shoot Apical Meristems of Oat (*Avena sativa* L.). *J. Plant Physiol.* 148, 667–671.
- Zhang, S., Cho, M.-J., Koprek, T., Yun, R., Bregitzer, P., Lemaux, P.G., 1999. Genetic transformation of commercial cultivars of oat (*Avena sativa* L.) and barley (*Hordeum vulgare* L.) using in vitro shoot meristematic cultures derived from germinated seedlings. *Plant Cell Rep.* 18, 959–966.
- Zhang, S., Williams-Carrier, R., Lemaux, P., 2002. Transformation of recalcitrant maize elite inbreds using in vitro shoot meristematic cultures induced from germinated seedlings. *Plant Cell Rep.* 21, 263–270.